

Section 24(c) Special Local Need Label

S-METOLACHLOR GROUP 15 HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS

Dual Magnum® Herbicide For Weed Control in Watermelon

EPA Reg. No. 100-816 EPA SLN No. TX-190005

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2030

SYNGENTA'S SPECIAL CONDITIONS, RISKS OF USE AND DISCLAIMER FOR USE OF DUAL MAGNUM HERBICIDE ON CROPS ON THIS 24(c) LABEL

IMPORTANT- READ BEFORE USE

THESE CONDITIONS RISKS OF USE AND DISCLAIMER ARE REQUIRED BY SYNGENTA CROP PROTECTION LLC AND NOT SPECIFIED BY U.S. EPA OR THE OF STATE OF TEXAS

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, SYNGENTA CROP PROTECTION, LLC INTENDS THAT THE PRODUCT THAT IS THE SUBJECT OF THIS SECTION 24(c) LABEL BE PURCHASED ONLY BY END USERS WHO AGREE BY ELECTRONIC SIGNATURE ON SYNGENTA CROP PROTECTION'S INTERNET SITE TO THE TERMS AND CONDITIONS REQUIRED BY SYNGENTA CROP PROTECTION, LLC INCLUDING A WAIVER AND RELEASE FROM ALL LIABILITY AND INDEMNIFICATION BY THE USER AND/OR GROWER OF SYNGENTA AND OTHERS FOR FAILURE TO PERFORM AND FOR CROP INJURY, CROP YIELD REDUCTION, AND/OR CROP LOSS FROM USE OF DUAL MAGNUM HERBICIDE ON CROPS ON THIS 24(c) LABEL. IF SUCH TERMS AND CONDITIONS ARE UNACCEPTABLE, RETURN THE DUAL MAGNUM HERBICIDE AT ONCE UNOPENED OR USE THE DUAL MAGNUM HERBICIDE FOR A DIFFERENT APPROVED USE IN ACCORDANCE WITH THE LABEL AFFIXED TO THE PRODUCT CONTAINER.

USE OF DUAL MAGNUM HERBICIDE (THE "PRODUCT") ON CROPS LISTED (THE "CROP") FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP. SYNGENTA RECOMMENDS THAT THE USER TEST THIS PRODUCT TO DETERMINE ITS SUITABILITY FOR SUCH INTENDED USE.

Syngenta Crop Protection, LLC makes the Product available for use in the manner described in this Section 24(c) Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crop outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss.

The decision to use this Product in the manner described in this Section 24(c) Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors. Syngenta recommends that the user test this Product to determine its suitability for such intended use.

By purchasing the Product for use, or using the Product in the manner described in this Section 24(c) Labeling, you acknowledge and accept that, to the extent consistent with applicable law:

- 1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- 2) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crop other than the statements on this Supplemental labeling;
- 3) Syngenta Crop Protection, LLC do not make, and do not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crop and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- 4) Syngenta Crop Protection, LLC disclaim all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- 5) these conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Syngenta Crop Protection, LLC or their respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.
- The risk of crop injury increases when adjuvants (e.g., non-ionic surfactants, crop oils, etc.), nitrogen sources (e.g., AMS, UAN), fertilizers or other pesticides are applied with Dual Magnum Herbicide.

WATERMELON

Dual Magnum Herbicide may be applied to direct seeded or transplanted crops for residual control or suppression of annual grasses, yellow nutsedge and small seeded broadleaf weeds.

Dual Magnum Herbicide will not control emerged weeds so use an appropriate registered foliar herbicide or mechanical or physical methods to control emerged weeds.

Application Timing	Crop Growth Stage	Rate (pt/A) ¹	Pre-harvest Interval ²
Preplant ³	Before planting	0.67 - 1.33	- 60 days
Pre-emergence	Before emergence or transplanting		
Post-emergence	After emergence or transplanting	0.67 - 1.33	
Post-emergence row middles			

¹ Use lower rates on coarse-textured soils and higher rates on fine-textured soils

Row Middle Application for Bareground: Apply Dual Magnum Herbicide at 0.5 to 0.75 pt/A between crop rows using a shielded or hooded sprayer. Spray contact with the crop or on the soil within 6 inches of the crop roots may lead to crop stunting.

Row Middle Application for Mulch: Apply Dual Magnum Herbicide at 0.5 to 0.75 pt/A between crop rows using a shielded or hooded sprayer. In raised bed mulch, spray can contact sides of bed covered in mulch but spray must not contact the top of the mulched bed or crop foliage. In flat bed mulch, avoid contact with top of the mulched bed as washing of the herbicide into the plant hole could be injurious. Spray contact with the crop or on the soil within 6 inches of the crop roots may lead to crop stunting.

If crops are grown on plastic mulch, the Dual Magnum Herbicide pre-emergence application should be made before laying the plastic. Dual Magnum Herbicide may also be applied as a row middle application after the laying of the plastic mulch.

² Time between Dual Magnum Herbicide application and crop harvest

³ No more than 14 days prior to planting

Dual Magnum Herbicide can be applied as part of a sequential weed control program. If Dual Magnum Herbicide was applied preplant, pre-emergence or pre-transplant, a second treatment of Dual Magnum Herbicide can be applied postemergence provided the total per crop does not exceed 1.33 pt/A.

Precautions:

- 1) If a pre-plant application is used, soil incorporation is not recommended. Minimize the mixing of herbicide treated soil into the seed furrow through the planting operations, so as to reduce the risk of crop injury.
- 2) The risk of post-emergence crop injury increases when adjuvants (e.g. non-ionic surfactants, crop oils, etc.), nitrogen sources (e.g. AMS, UAN), fertilizers or other pesticides are applied with Dual Magnum Herbicide.
- 3) All watermelon cultivars have not been studied for tolerance. Experiment on a limited basis until on-farm confidence in these use patterns and rates are obtained.

Restrictions:

- 1) Do not apply more than 1.33 pt/A (1.27 lb ai/A) of Dual Magnum Herbicide per year. Do not apply more than 1.27 lb ai/A per year of S-metolachlor-containing products.
- 2) Do not harvest for 60 days after the application of Dual Magnum Herbicide.

©2025 Syngenta

Dual Magnum® and the Syngenta logo are trademarks of a Syngenta Group Company

24(c) Registrant: Syngenta Crop Protection, LLC P. O. Box 18300 Greensboro, NC 27419-8300

Label Code: TX0816176CA0325