




For Agricultural or Commercial Use Only
EPA REG. No. 279-3158

EMERGENCY CALLS: 800-331-3148

EPA SLN NO.
TX-190007



FOR DISTRIBUTION AND USE ONLY IN TEXAS

For distribution and use only in the Texas counties of: Austin, Brazoria, Calhoun, Chambers, Colorado, Fort Bend (South and West of Hwy 36), Galveston, Jackson, Jefferson, Lavaca, Liberty, Matagorda, Victoria, Waller, and Wharton.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING

This label for Command® 3ME Microencapsulated Herbicide is valid until December 31, 2029 or until otherwise amended, withdrawn, canceled or suspended.

Section 24(c) State Registration

DIRECTIONS FOR USE:

For aerial application of Command® 3ME Microencapsulated Herbicide on rice.

Apply in a minimum of 5 gallons of finished spray per acre.

PRE-EMERGENT SURFACE BROADCAST APPLICATIONS

Command 3ME Microencapsulated Herbicide may be applied as a surface broadcast application 14 days prior to planting or up to 7 days after planting, but prior to weed emergence at the rate of 10.7 to 34.1 fl. oz. product (0.25 to 0.8 lb. active) per acre depending upon the soil texture. Refer to table below for specific rates and weeds controlled. For heavy soils use the higher specified rate, otherwise less than desirable weed control may result.

EARLY POSTEMERGENCE APPLICATIONS

Command 3ME Microencapsulated Herbicide may be applied after planting as an early post-emergence treatment to rice through the five-leaf stage to provide pre-emergence and residual control of grass weeds. Refer to table below for specific rates and weeds controlled. For control of existing grass present at the time of application include a post-emergence herbicide registered for the control of grass species in rice. Consult post-emergence herbicide label for specific directions regarding use rates and stage of weeds and crop.

Observe all buffer restrictions noted in the Restrictions section.

Partial weed control may result if levees are formed after Command 3ME Microencapsulated Herbicide has been applied.

Additional use of labeled post-emergent herbicide applications may be required where existing grass weeds are present at the time of application.

CROP	PESTS	RATE OF APPLICATION			
		Soil Texture	Broadcast Rates Per Acre*	Pounds active ingredient per acre	Acres per Gallon
Rice	Barnyardgrass (Watergrass) (Including herbicide resistant types) Broadleaf Signalgrass Crabgrass (Large, Smooth) Panicum (Common, Fall, Texas) Sprangletop	Coarse (light) Soils: (sand, loamy sand, sandy loam):	(10.7 – 14.1 fl oz)	0.25 - 0.33	11.6 – 9.1
		Medium Soils: (loam, silt, silt loam, sandy clay, sandy clay loam)	(17.1 – 21.3 fl oz)	0.4 - 0.5	7.5 - 6
		Fine (heavy) Soils: (silty clay, clay loam, silty clay loam, clay)	(21.3 – 34.1 fl oz)	0.5 - 0.8	6.0 – 3.76
* Select lower to higher rates based on lighter to heavier soil types.					

CROP SAFETY PRECAUTIONS:

Application of Command 3ME Microencapsulated Herbicide to fields that have been precision leveled with deep cuts may result in rice crop injury including stand loss. Consult with rice specialists for soil amending practices that can reduce potential for herbicide injury in precision- leveled fields.

MANDATORY SPRAY DRIFT MANAGEMENT:**Aerial Applications**

Aircraft used to apply Command 3ME Microencapsulated Herbicide shall be configured and operated in such manner as to minimize off-site spray movement to desirable species.

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to use a Coarse or coarser droplet size (ASABE S572.1).
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Aircraft

- Adjust Nozzles - Follow nozzle manufacturers recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

RELEASE HEIGHT - Aircraft

Higher release heights increase the potential for spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift

PRECAUTIONS:

Caution must be taken to minimize off target drift of Command 3ME Microencapsulated Herbicide as off-site movement of Command 3ME Microencapsulated Herbicide can cause foliar whitening or yellowing of some plants.

RESTRICTIONS:

Do not apply within 300 feet of downwind crops and other desirable non-target plants.

Do not apply within 1,200 feet of Towns and Housing Developments, Commercial Fruit, Nut, or Vegetable Production¹, Commercial Greenhouses or Nurseries.

¹ Except for vegetable crops registered for use on the Command 3ME Microencapsulated Herbicide label (including peppers, pumpkins, succulent peas, sweet corn, sweet potato and winter squash).

Command 3ME Microencapsulated Herbicide may be applied to water seeded rice from 14 days up to the pinpoint flood but prior to planting or during pegging up to re-flooding but prior to grass emergence.

With split applications do not apply more than a total of 34.1 fl oz/A Command 3ME Microencapsulated Herbicide (0.8 lb ai/A) per season.

Do not apply Command 3ME Microencapsulated Herbicide on rice fields in which concurrent crayfish or catfish farming is included in the cultural practices.

Do not use water containing Command 3ME Microencapsulated Herbicide residues from rice cultivation to irrigate food or feed crops, which are not registered for use with Command 3ME Microencapsulated Herbicide.

For distribution and use only in the Texas counties of:

Austin, Brazoria, Calhoun, Chambers, Colorado, Fort Bend (ONLY South and West of Hwy. 36), Galveston, Jackson, Jefferson, Lavaca, Liberty, Matagorda, Victoria, Waller, and Wharton.

"This product should only be applied when the potential for drift to adjacent sensitive areas (known habitat for endangered species) is minimal (wind is blowing away from sensitive area)".

REPLANTING INSTRUCTIONS

If initial planting of rice fails to produce a uniform stand, rice may be replanted in fields treated with Command 3ME Microencapsulated Herbicide. DO NOT retreat fields with a second pre-plant application of Command 3ME Microencapsulated Herbicide. DO NOT replant treated fields with any crop at intervals that are inconsistent with the ROTATIONAL CROP GUIDELINES on this labeling.

ROTATIONAL CROP GUIDELINES

Rotate to crops as listed below, otherwise crop injury may occur.

Note: When using Command 3ME Microencapsulated Herbicide with other registered herbicides always refer to rotational restrictions and precautions on the other product's label.

ANYTIME - Cotton*, Peas, Peppers, Pumpkins, Rice, Soybeans, Tobacco

9 MONTHS - Cotton, Corn (Field, Pop, Seed, Sweet), Cucurbits, Dry Beans, Peanuts, Potatoes, Snap Beans, Sorghum, Sugar Beets, Sweet Potatoes, Tomatoes (Transplanted)

12 MONTHS - All crops

- Refer to "REQUIREMENTS FOR PLANTING TIME APPLICATIONS" and "REPLANTING INSTRUCTIONS" in the Command 3ME Section 3 label.

ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

Conditions of Sale and Limitation of Warranty and Liability:

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions beyond the control of FMC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold FMC and Seller harmless for any claims relating to such factors.

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the Directions for Use when used in accordance with the directions under normal conditions of use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, FMC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE, NOR ANY OTHER EXPRESS OR IMPLIED WARRANTIES WITH RESPECT TO THE SELECTION, PURCHASE, OR USE OF THIS PRODUCT. Any warranties, express or implied, having been made are inapplicable if this product has been contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to (or beyond the control of) Seller or FMC, and Buyer assumes the risk of any such use.

To the extent consistent with applicable law, FMC or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF FMC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF FMC OR SELLER, THE REPLACEMENT OF THE PRODUCT.

This Conditions of Sale and Limitation of Warranty and Liability may not be amended by any oral or written agreement.

NOTICE TO BUYER -- Purchase of this material does not confer any rights under patents of countries outside of the United States.

FMC, the FMC logo, and Command are trademarks of FMC Corporation and/or an affiliate.

© 2024 FMC Corporation All rights reserved.

Label Code : R-4894 101124 09-18-24