



ITERA

Section 24(c) Special Local Need Label

S-METOLACHLOR	GROUP	15	HERBICIDE
METRIBUZIN	GROUP	5	HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS

Boundary® 6.5 EC Herbicide

FOR CONTROL OF GLYPHOSATE-RESISTANT ITALIAN RYEGRASS IN THE FALL PRIOR TO SOYBEAN OR CORN PLANTING THE FOLLOWING SPRING

**EPA Reg. No. 100-1162
EPA SLN No. TX-200005**

This label for Boundary 6.5EC Herbicide expires and must not be distributed or used in accordance with this SLN registration after December 31, 2030.

KEEP OUT OF REACH OF CHILDREN CAUTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

DIRECTIONS FOR USE

FALL APPLICATION FOR CONTROL OF GLYPHOSATE- RESISTANT ITALIAN RYEGRASS (*Lolium multiflorum*) PRIOR TO SOYBEAN OR CORN PLANTING

Apply Boundary® 6.5EC at 2.0 pt/A (1.62 lb ai/A) in the fall (September 1 – November 30) when rotating to soybean or corn. If applied after November 30, apply Boundary 6.5EC at 1.6 pt/A (1.31 lb ai ai/A) when rotating to corn. Regardless of the application date, do not apply Boundary 6.5EC to frozen ground.

A tillage operation may precede the Boundary 6.5EC application. A fall and/or spring tillage operation may follow application but do not exceed an incorporation depth of 2-3 inches.

Gramoxone® SL 3.0 at 1.3 to 2.7 pt/A can be tank mixed with Boundary 6.5EC to control emerged Italian ryegrass. Add a nonionic surfactant at 1 quart/100 gallons (0.25% v/v) in mixtures with Gramoxone SL 3.0. Read and follow all directions and restrictions on the Gramoxone SL 3.0 label.

Crop Rotation Intervals^{1,3}

Rotational Interval After Application of Boundary 6.5EC ²				
1 Month	4½ Months	8 Months	12 Months	18 Months
Corn ³	Winter Barley Winter Wheat Alfalfa	Peas Rice Spring Barley Spring Wheat	Asparagus Cotton Forage Grasses Lentils Sainfoin Sugarcane Tomatoes Other Crops not listed (except root crops)	Onions Sugar Beets and Other Root Crops

¹Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed. Stand reductions may occur in some areas.

²Crop rotation recommendations do not include restrictions for the tank mix partner. Refer to the label of the other product for additional restrictions.

³The 1 month rotational restriction only applies when Boundary 6.5EC is applied at 1.6 pt/A (1.31 lb ai/A). Corn rotation is 4 months for rates greater than 1.6 pt/A Boundary 6.5EC.

Restrictions

- Do not make more than one fall application of Boundary 6.5EC for control of Italian ryegrass.
- Do not apply more than 2 pt/A (1.62 lb ai/A) of Boundary 6.5EC in a single application prior to November 30 if rotating to soybeans or corn the following spring.
- The total amount of Boundary 6.5EC applied (fall + spring) cannot exceed 3.9 pints/A/year (2.56 lb of S-metolachlor/A/year and 0.61 lb of metribuzin/A/year).
- Do not use sand, loamy sand, or sandy soils with less than 0.5% organic matter or soils with pH greater than 7.0, if rotating to corn following application after November 30.
- Do not exceed 3.71 lb ai/A/year of S-metolachlor-containing products. Do not exceed 1.0 lb ai/A/year of metribuzin-containing products.

Boundary® and Gramoxone® are trademarks of a Syngenta Group Company

©2025 Syngenta

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300

Label Code: TX1162193ABA0825