

**RESTRICTED USE PESTICIDE  
DUE TO ACUTE TOXICITY  
FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS– NOT TO BE  
USED BY UNCERTIFIED PERSONS WORKING UNDER THE SUPERVISION OF A  
CERTIFIED APPLICATOR**

**syngenta.**

**INTRA**

## **Section 24(c) Special Local Need Label**

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS**

**Gramoxone® SL 3.0  
Harvest Aid Applications on Cotton in Texas  
(Except in the Counties of Starr, Hidalgo, Willacy, and Cameron)**

**EPA Reg. No. 100-1652  
EPA SLN No. TX-240003**

**This label expires and must not be distributed or used in accordance with this SLN  
registration after December 31, 2028**

**KEEP OUT OF REACH OF CHILDREN / MANTENER FUERA DEL ALCANCE DE LOS  
NIÑOS.**



**DANGER / POISON  
PELIGRO / VENENO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### **DIRECTIONS FOR USE**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered label.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL  
MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

## Specific Crop Use Directions

Three multiple broadcast harvest aid applications of Gramoxone SL 3.0 may be used. **Do not exceed a total of 1.3 pt/A/year of Gramoxone SL 3.0 or 0.50 lb ai/A/year of paraquat containing products for harvest aid uses.** Use a minimum total spray volume per acre of 10 gallons by ground, and 5 gallons by air.

Crop	Rate per Acre	Precautions, Restrictions and Comments
Tank mixes with phosphate or chlorate defoliant	5.4-10.7 fl oz alone <u>or</u> tank mixed with 1 1/2 - 2 gal chlorate, <u>or</u> 1 1/2 - 2 pt phosphate	<ul style="list-style-type: none"> <li>• Do not harvest within 7 days of application when used with phosphate or chlorate defoliant.</li> <li>• In desert cotton areas or on rank vigorous cotton, use the 10.7 fl oz rate of Gramoxone SL 3.0</li> <li>• Apply when 75% or more of the bolls are open and remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited.</li> <li>• Follow Harvest Aid Use Precautions listed on the Federal label.</li> </ul>
Tank mixes for boll opening and defoliation of southern cotton	2.7-3.3 fl oz	<ul style="list-style-type: none"> <li>• To aid in defoliation and opening of mature bolls, Gramoxone SL 3.0 may be tank mixed with the following products: DEF<sup>®</sup> defoliant, DROPP<sup>®</sup> defoliant, PREP<sup>™</sup> PGR, FOLEX<sup>®</sup> defoliant.</li> <li>• Apply when 60% or more of the bolls are open and the remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited.</li> <li>• Refer to tank mix product label for rate, directions, limitations and caution</li> </ul>
Harvest aid for defoliation and boll opening of stripper or spindle harvested cotton	2.1-7.5 fl oz	<ul style="list-style-type: none"> <li>• Apply when 75% of the bolls are open and the remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited. Slice bolls and inspect the seed for maturity.</li> <li>• Gramoxone SL 3.0 may be applied alone or tank mixed with the cotton harvest aid products listed above.</li> <li>• To avoid leaf sticking, apply Gramoxone SL 3.0 as a desiccant approximately 3-7 days after defoliation or a conditioning application and 7-14 days before harvest.</li> <li>• Cooler temperatures may cause a longer waiting period between application of Gramoxone SL 3.0 as a desiccant and defoliation/conditioning.</li> <li>• Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation.</li> </ul>
Harvest aid desiccation of stripper cotton or spindle harvested cotton late season	11.2-21 fl oz	<ul style="list-style-type: none"> <li>• Use to desiccate regrowth occurring after defoliation or desiccation.</li> <li>• Apply when 85% of the bolls are open and the remaining bolls to be harvested are mature.</li> <li>• Development of immature bolls will be inhibited. Slice bolls and inspect the seed for maturity.</li> <li>• To avoid leaf sticking, apply Gramoxone SL 3.0 as a desiccant approximately 3-7 days after defoliation or a conditioning application and 7-14</li> </ul>

Crop	Rate per Acre	Precautions, Restrictions and Comments
		days before harvest. <ul style="list-style-type: none"> <li>• Lower rates in the range may be necessary south of I-10 in Texas where temperatures are typically higher during defoliation.</li> </ul>
Desiccation of Regrowth	11.2-21 fl oz	<ul style="list-style-type: none"> <li>• Use to desiccate regrowth after defoliation or desiccation.</li> <li>• Regrowth is difficult to control, therefore, thorough coverage with the full recommended rate is necessary.</li> <li>• Control is dependent upon growing condition and desiccation of small new growth may not always be complete.</li> <li>• Use higher rate if regrowth is excessive</li> </ul>

Refer to the Gramoxone SL 3.0 federal label for additional cotton recommended uses and nonionic surfactant recommendations.

**Additional Precautions, Restrictions and Comments:**

- It is advisable, because of extremes in environmental and plant conditions, to apply the range of rates on a small block of cotton to determine the rate that best fits your needs.
- Do not pasture lactating dairy animals.
- Do not pasture livestock in treated fields or feed treated foliage.
- When combined with phosphate defoliant, observe livestock cautions listed on the phosphate label.
- Do not harvest within 7 days of application when used with phosphate or chlorate defoliant.
- Do not apply this product through any type of irrigation system

©2023 Syngenta

Gramoxone® and the Syngenta logo are trademarks of a Syngenta Group Company  
 Def®, Dropp®, Prep™ PGR, are trademarks of Bayer CropScience  
 Folex® is a trademark of Amvac Chemical Corporation

24(c) Registrant:  
 Syngenta Crop Protection, LLC  
 P. O. Box 18300  
 Greensboro, NC 27419-8300

Label Code: TX1652204AA0921