



RESTRICTED USE PESTICIDE

Due to Toxicity to Fish and Aquatic Organisms.
For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

BETA-CYFLUTHRIN GROUP 3 INSECTICIDE

FIFRA Section 24(c) Special Local Need (SLN) Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS

For Control of Asian Citrus Psyllid on Citrus.

BAYTHROID® XL

EPA REG No. 264-840

EPA SLN No. TX-260002

**THIS LABEL IS VALID UNTIL DECEMBER 31, 2030, UNLESS OTHERWISE AMENDED,
WITHDRAWN, CANCELED, OR SUSPENDED.**

ACTIVE INGREDIENT:

β-cyfluthrin

Cyano(4-fluoro-3-phenoxyphenyl)methyl-3-(2,2-dichloroethenyl)-2,2-dimethyl-cyclopropanecarboxylate 12.70%

OTHER INGREDIENTS: 87.30%

TOTAL: 100.00%

**Contains 1 lb Beta-cyfluthrin per gallon.
(This product contains aromatic petroleum distillates.)**

DIRECTIONS FOR USE

Restricted Use Pesticide

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and the federal label for this product must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the EPA registered label.

TREE and VINE CROPS CROP USE DIRECTIONS

For all crops, apply specific dosage of BAYTHROID XL at early threshold for target pest, as population begins to develop. Degree of control or suppression of additional labeled pests will be determined, in part by the stage of pest development at application and infestation level of those pests.

Specified application rates within this label are based on full-size mature trees and vines. Application timing should be based on careful scouting and local economic thresholds. Use the higher rates for moderate to heavy insect pressure or when applying by air. Lower rates are generally adequate for smaller trees/vines or low to moderate insect pressure but require careful scouting and may require more frequent application.

BAYTHROID XL is an Emulsifiable Concentrate (EC) formulation and is active by contact and ingestion. For tree and vine crops, apply by ground or air equipment using sufficient water to obtain through coverage of target plant parts for optimum performance.

CITRUS

Includes all members of Crop Group 10:

Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Orange (sweet and sour), Pummelo, Satsuma mandarin, White sapote, and other cultivars and/or hybrids of these.

PESTS CONTROLLED	Rate fluid oz/Acre	Rate lb AI/Acre
Asian Citrus Psyllid	3.2 - 6.4	0.025 - 0.05

Foliar Application Restrictions

Pre-Harvest Interval (PHI): **0 day**.

Maximum BAYTHROID XL allowed per 7-day interval: **6.4 fluid oz/A (0.05 lb AI/Acre)**.

Maximum BAYTHROID XL allowed per crop season: **6.4 fluid oz/A (0.05 lb AI/Acre)**.

Minimum application volume (water): 25 GPA – ground, 25 GPA – aerial application.

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

*As with any crop-protection product, always read and follow label directions.
For additional information call toll-free 1-866-99BAYER (1-866-992-2937).*

**Bayer CropScience LLC
800 N. Lindbergh Blvd.
St. Louis, MO 63167
1-866-99BAYER (1-866-992-2937)**